


## Corsodyl Mouthwash

### *Instructions for use:*

1. Use 10ml (up to the line in the bottle lid)
1. Twice daily
2. Rinsing around the mouth for 1 minute
3. Use for 2 weeks
4. Then use for 1 week in every month if needed.



Most tooth pastes contain a foaming agent (called SLS) that interferes with the affect of Corsodyl.

If using the mouthwash at the same time as brushing then it is important to use a tooth paste free of SLS, such as:

- **Corsodyl Daily gum and tooth paste**
- **Sensodyne Pro Enamel**

It is otherwise recommended that Corsodyl is used between 30 minutes to 2hrs after brushing with a tooth paste containing SLS so that the mouthwash is not inhibited by any paste still in the mouth.

N. PRASAD, B.D.S. (HONS), M.F.D.S. R.C.S. (ENG), R. V. MARSDEN, B.D.S.,  
L. FROST, B.D.S., F.D.S. R.C.P.S. (GLAS) SPECIALIST IN ORAL SURGERY

THE RED HOUSE DENTAL PRACTICE  
VICTORIA ROAD, MALTON, NORTH YORKSHIRE YO17 7JJ.  
TEL. 01653 693809  
EMAIL: [reception@redhousedentists.co.uk](mailto:reception@redhousedentists.co.uk)  
[www.redhousedentists.co.uk](http://www.redhousedentists.co.uk)