



Tooth Whitening

- **Before bed**
 - ✓ **Brush teeth** thoroughly with a sensitive toothpaste.
 - ✓ **Rinse with water.**
 - ✓ **Prepare your tray:** Place a little whitening gel into each of the reservoirs.
 - ✓ **Place the tray in the mouth,** and gently ensure it is well seated.
 - ✓ **Remove excess whitening gel** with a cotton bud, tissue, or toothbrush.
 - ✓ **Do not swallow any excess material.** Spit it out.
- **On waking**
 - ✓ **Remove the tray from your mouth.**
 - ✓ **Rinse your mouth with water.**
 - ✓ **Brush with sensitive toothpaste.**
 - ✓ **Clean the trays with cool water,** using a cotton bud or toothbrush to help.
 - ✓ **Rinse and clean the tray.**
 - ✓ **Leave tray in the open air to dry**

Whitening for sensitive teeth:

- **If your teeth are already sensitive before treatment:**
 - ✓ **For 1 week before whitening,** place a sensitive toothpaste (eg. Colgate Pro Relief/Sensodyne Pro Enamel) in the bleaching tray each evening for 1 to 2 hours.
 - ✓ Brush with a sensitive toothpaste twice daily.
 - ✓ When whitening the teeth begins, wear the tray during the day for 2 hours each day only for the first week, and then nightly for the following weeks.
- **If sensitivity arises during treatment:**
 - ✓ **Place a sensitive toothpaste in the tray, wear overnight.**
 - ✓ On waking, brush your teeth with a sensitive toothpaste.
 - ✓ Continue with your normal whitening regime.
- **If the teeth are frequently sensitive:**
 - ✓ **Wear the tray loaded with sensitive toothpaste on alternate nights to the whitening gel,** ie. place sensitive toothpaste/fluoride gel in the tray one night, and tooth whitening gel the next.
 - ✓ On the nights you are due to whiten your teeth, also wear the tray loaded with a sensitive toothpaste for *30 minutes before whitening, and 30 minutes after.*
 - ✓ Contact your dentist if you have any concerns.

Other tips for sensitive teeth:

- ✓ If the teeth become too sensitive to continue, **don't be afraid to stop treatment for a few days** to give the teeth a rest! Apply a sensitive toothpaste (Colgate Pro Relief/Sensodyne Pro Enamel)
- ✓ **Reduce the treatment time you wear the tray; wear the tray less often.**
- ✓ Do not overload the tray, and use slightly less gel in the tray.

N. PRASAD, B.D.S. (HONS), M.F.D.S. R.C.S. (ENG), R. V. MARSDEN, B.D.S.,
L. FROST, B.D.S., F.D.S. R.C.P.S. (GLAS) SPECIALIST IN ORAL SURGERY

THE RED HOUSE DENTAL PRACTICE
VICTORIA ROAD, MALTON, NORTH YORKSHIRE YO17 7JJ.
TEL. 01653 693809
EMAIL: reception@redhousedentists.co.uk
www.redhousedentists.co.uk

